

A flow platform for degradation-free CuAAC bioconjugation

Citation for published version:

Hatit, MZC, Reichenbach, LF, Tobin, JM, Vilela, F, Burley, GA & Watson, AJB 2018, 'A flow platform for degradation-free CuAAC bioconjugation', *Nature Communications*, vol. 9, 4021.
<https://doi.org/10.1038/s41467-018-06551-0>

Digital Object Identifier (DOI):

[10.1038/s41467-018-06551-0](https://doi.org/10.1038/s41467-018-06551-0)

Link:

[Link to publication record in Heriot-Watt Research Portal](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Nature Communications

Publisher Rights Statement:

© The Author(s) 2018. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

General rights

Copyright for the publications made accessible via Heriot-Watt Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

Heriot-Watt University has made every reasonable effort to ensure that the content in Heriot-Watt Research Portal complies with UK legislation. If you believe that the public display of this file breaches copyright please contact open.access@hw.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

ARTICLE

DOI: 10.1038/s41467-018-06551-0

OPEN

A flow platform for degradation-free CuAAC bioconjugation

Marine Z.C. Hatit¹, Linus F. Reichenbach¹, John M. Tobin², Filipe Vilela²,
Glenn A. Burley¹ & Allan J.B. Watson³

The Cu-catalyzed azide-alkyne cycloaddition (CuAAC) reaction is a cornerstone method for the ligation of biomolecules. However, undesired Cu-mediated oxidation and Cu-contamination in bioconjugates limits biomedical utility. Here, we report a generic CuAAC flow platform for the rapid, robust, and broad-spectrum formation of discrete triazole bioconjugates. This process leverages an engineering problem to chemical advantage: solvent-mediated Cu pipe erosion generates ppm levels of Cu in situ under laminar flow conditions. This is sufficient to catalyze the CuAAC reaction of small molecule alkynes and azides, fluorophores, marketed drug molecules, peptides, DNA, and therapeutic oligonucleotides. This flow approach, not replicated in batch, operates at ambient temperature and pressure, requires short residence times, avoids oxidation of sensitive functional groups, and produces products with very low ppm Cu contamination.

¹Department of Pure and Applied Chemistry, University of Strathclyde, 295 Cathedral Street, Glasgow G1 1XL, UK. ²Chemical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK. ³School of Chemistry, University of St Andrews, North Haugh, St Andrews KY16 9ST, UK. Correspondence and requests for materials should be addressed to G.A.B. (email: glenn.burley@strath.ac.uk) or to A.J.B.W. (email: aw260@st-andrews.ac.uk)




Fig. 2 Development of a flow CuAAC process based on H₂O Cu erosion. **a** The effect of H₂O on the CuAAC reaction efficiency using three representative alkynes; **b** Correlation of [Cu] vs. solvent composition ((H₂O)); **c** Demonstration of the flow effect—14 ppm Cu CuAAC reactions in flow and in flask

be very low; the increased circulation established under the flow set up would enhance mass transport to provide reaction efficiency not possible in batch²⁶.

This hypothesis was found to be valid. Three benchmark CuAAC reactions, using three alkynes (**1a–1c**), with known differences in reactivity with benzyl azide (**2**) were evaluated in a laminar flow system comprising a pump and copper reactor at ambient temperature and pressure (Fig. 2a). The reaction does not proceed in pure MeCN or pure H₂O and [Cu] in the eluent (10 mL collected, 1 mL/min under ambient conditions) was extremely low (<20 ppm). However, the addition of small amounts of H₂O to the bulk MeCN resulted in the formation of triazole **3a–c**, which peaked at 5:1 solvent mixture. Control experiments with an unused Cu reactor (Fig. 2b, red bars) vs. a reactor used for CuAAC reactions (Fig. 2b, blue bars) demonstrated greater erosion in the used reactor, consistent with a more exposed surface due to repeated chemistry; however the solvent composition/erosion trend was comparable, peaking at 1:1 H₂O/MeCN. The addition of small percentages of H₂O to the carrier solvent (MeCN) enabled the CuAAC reaction of equimolar ynamine **1a** and BnN₃ (**2**) effectively at 5:1 MeCN:H₂O (t_R = 10 min; Fig. 2a). Whilst ynamine **1a** exhibits faster batch-reaction kinetics based on a pK_a modifying Cu-ligation⁴², the mixed solvent system was also effective at enabling the CuAAC reaction of more standard alkynes **1b** and **1c** at the same flow rate. Analysis of the eluent by ICP-MS revealed that [Cu] was ~14 ppm, which is well below the limit required for use in vivo applications^{5,11}. Importantly, control experiments identified a flow phenomenon. Attempting the CuAAC reaction of alkynes **1a–1c** in flask experiments at 14 ppm Cu was unsuccessful for **1b** and **1c** and only moderately successful for the more reactive ynamine **1a** (53% yield after 72 h), whereas the flow system results in quantitative conversion in 10 min (Fig. 2c). Residence times were also shortened significantly to ca. 1 min for more reactive substrates.

Scope of the flow platform. The scope of the flow CuAAC process was both broad and reproducible using three different alkyne classes (**1a–1c**) across a series of azide substrates (**3–20**; Fig. 3). Triazole products derived from simple azides, azido fluorophores, and azide possessing specific functions for downstream applications, were all isolated in high yield after a single pass. Importantly, ICP-MS analysis of the products again found the residual [Cu] was <20 ppm (see Supporting Information for full details).

We also examined the compatibility of the flow process with regards to established CuAAC chemoselectivity profiles (Fig. 3). Dityne **18**, containing aliphatic alkyne and aromatic ynamine sites, underwent sequential CuAAC ligation, firstly with the coumarin azide **19** at the ynamine site followed by ligation with the nucleobase azide **20** at the aliphatic alkyne site; complete chemoselectivity was observed throughout. This demonstrates that established reactivity profiles⁴³ are replicated in the flow format and that our system enhances not only overall reaction kinetics but does so at very low [Cu].

The biomedical utility of the CuAAC reaction lies primarily in the ligation of bio-relevant molecules. We assessed the flow CuAAC process as a method for the ligation of representative alkyne-derivatives of nucleic acids and peptides, which have known susceptibility to form oxidized byproducts in the presence of a Cu catalyst (Fig. 4)^{8,10}. Installation of a fluorinated residue onto a marketed PARP inhibitor⁴⁴, and a common fluorophore onto a series of peptides and DNA strands containing oxidizable functionality produced triazole products with minimal formation of side-products. These include CuAAC ligations with oligodeoxyribonucleotides (ODNs) and the core ApolipoproteinE (ApoE) peptide sequence (**27**)⁴⁵, which has demonstrated utility as a delivery vehicle across the blood brain barrier⁴⁶. Residues with known oxidative susceptibility (**27a–e**) under conventional CuAAC batch conditions were installed on the N-terminus to report any potential degradation




Fig. 3 Scope of the flow CuAAC process. For each product number (in bold), data are reported as percent isolated yield. Products with designation **a** are derived from alkyne **1a**, **b** from alkyne **1b**, and **c** from alkyne **1c**

by reactive oxygen species and formed the expected triazole products (1 mL/min; t_R = 8 min), with trace Cu contamination and no associated degradation.

Bioconjugation. Finally, we explored applying our flow-based CuAAC ligation approach to prepare therapeutic bioconjugates. Phosphoramidate morpholino oligonucleotides (PMOs) are a class of oligonucleotides with established therapeutic importance^{47–49}. An essential requirement for in vivo efficacy of this class of biologics is the need conjugate a cell penetrating peptide sequence onto one of the termini to enable effective delivery to the central nervous system. The bioconjugate triazole **28** was prepared from precursors derived from a PMO azide with known

in vivo efficacy as a splice-switching oligonucleotide for the treatment of Spinal Muscular Atrophy (SMA) and a peptide fragment derived from a portion of the ApoE protein⁵⁰. Under flow conditions, the ApoE-PMO bioconjugate (**28**) was formed in 60% yield after 15 passes (1 mL/min; total t_R = 30 min). No reaction was observed after 24 h under equivalent batch conditions, with only 26% yield of **28** obtained in batch after 48 h using 100 equiv Cu.

Discussion

In summary, we have developed a rapid and operationally simple flow-based platform for the CuAAC reaction that operates at ambient temperature and pressure. Solvent-induced erosion of a




Fig. 4 Alkyne scope of the flow CuAAC process. For each product number (in bold), data are reported as percent isolated yield

Cu pipe provides catalytically competent Cu to promote the CuAAC reaction of a range of both small molecules and bio-molecules without oxidative damage to labile functional groups and with trace Cu contamination. We have demonstrated the dependency of the system on the composition of the medium and that the observed effect is unique to the flow conditions with comparable isolated experiments of low efficiency. We expect that these findings will significantly increase the utility of flow-assisted CuAAC across a series of academic and industrial applications.

Methods

General methods. See Supplementary Methods for further details supporting experiments, Supplementary Tables 1–11 for additional data, and Supplementary Figures 1–125 for spectra.

Data availability

All data generated or analyzed during this study are included in this published article (and its supplementary information files). These data are also available from the author upon request.

Received: 29 August 2018 Accepted: 11 September 2018

Published online: 01 October 2018

References

- Kolb, H. C., Finn, M. G. & Sharpless, K. B. Click chemistry: diverse chemical function from a few good reactions. *Angew. Chem. Int. Ed.* **40**, 2004–2021 (2001).
- Hein, J. E. & Fokin, V. V. Copper-catalyzed azide–alkyne cycloaddition (CuAAC) and beyond: new reactivity of copper(I) acetylides. *Chem. Soc. Rev.* **39**, 1302–1315 (2010).
- Meldal, M. & Tornøe, C. W. Cu-catalyzed azide–alkyne cycloaddition. *Chem. Rev.* **108**, 2952–3015 (2008).
- Moses, J. E. & Moorhouse, A. D. The growing applications of click chemistry. *Chem. Soc. Rev.* **36**, 1249–1262 (2007).
- Sletten, E. M. & Bertozzi, C. R. Bioorthogonal chemistry: fishing for selectivity in a sea of functionality. *Angew. Chem. Int. Ed.* **48**, 6974–6998 (2009).
- McKay, C. S. & Finn, M. G. Click chemistry in complex mixtures: bioorthogonal bioconjugation. *J. Chem. Biol.* **21**, 1075–1101 (2014).
- Saito, F., Noda, H. & Bode, J. W. Critical evaluation and rate constants of chemoselective ligation reactions for stoichiometric conjugations in water. *ACS Chem. Biol.* **10**, 1026–1033 (2015).
- Gierlich, J., Burley, G. A., Gramlich, P. M. E., Hammond, D. M. & Carell, T. Click chemistry as a reliable method for the high-density postsynthetic functionalization of alkyne-modified DNA. *Org. Lett.* **8**, 3639–3642 (2006).
- Hong, V., Presolski, S. I., Ma, C. & Finn, M. G. Analysis and optimization of copper-catalyzed azide–alkyne cycloaddition for bioconjugation. *Angew. Chem. Int. Ed.* **48**, 9879–9883 (2009).
- Li, S. et al. Extent of the oxidative side reactions to peptides and proteins during the CuAAC reaction. *Bioconjugate Chem.* **27**, 2315–2322 (2016).
- Abel, G. R. Jr., Calabrese, Z. A., Ayco, J., Hein, J. E. & Ye, T. Measuring and suppressing the oxidative damage to DNA during Cu(I)-catalyzed azide–alkyne cycloaddition. *Bioconjugate Chem.* **27**, 698–704 (2016).
- Dommerholt, J., Rutjes, F. P. J. T. & van Delft, F. L. Strain-promoted 1,3-dipolar cycloaddition of cycloalkynes and organic azides. *Top. Curr. Chem.* **374**, 16 (2016).
- Knall, A.-C. & Slugovc, C. Inverse electron demand diels–alder (iEDDA)-initiated conjugation: a (high) potential click chemistry scheme. *Chem. Soc. Rev.* **42**, 5131–5142 (2013).
- Lang, K. & Chin, J. W. Bioorthogonal reactions for labeling proteins. *ACS Chem. Biol.* **9**, 16–20 (2014).
- Baskin, J. M. et al. Copper-free click chemistry for dynamic in vivo imaging. *Proc. Natl Acad. Sci. USA* **104**, 16793–16797 (2007).
- Li, X., Fang, T. & Boons, G.-J. Preparation of well-defined antibody–drug conjugates through glycan remodeling and strain-promoted azide–alkyne cycloadditions. *Angew. Chem. Int. Ed.* **53**, 7179–7182 (2014).
- Blackman, M. L., Rozyen, M. & Fox, J. M. Tetrazine ligation: fast bioconjugation based on inverse-electron-demand diels–alder reactivity. *J. Am. Chem. Soc.* **130**, 13518–13519 (2008).
- Taylor, M. T., Blackman, M. L., Dmitrenko, O. & Fox, J. M. Design and synthesis of highly reactive dienophiles for the tetrazine–trans-cyclooctene ligation. *J. Am. Chem. Soc.* **133**, 9646–9649 (2011).
- Devaraj, N. K., Upadhyay, R., Haun, J. B., Hilderbrand, S. A. & Weissleder, R. Fast and sensitive pretargeted labeling of cancer cells through a tetrazine/trans-cyclooctene cycloaddition. *Angew. Chem. Int. Ed.* **48**, 7013–7016 (2009).
- Devaraj, N. K., Weissleder, R. & Hilderbrand, S. A. Tetrazine-based cycloadditions: application to pretargeted live cell imaging. *Bioconjugate Chem.* **19**, 2297–2299 (2008).
- Murrey, H. E. et al. Systematic evaluation of bioorthogonal reactions in live cells with clickable HaloTag Ligands: Implications for Intracellular Imaging. *J. Am. Chem. Soc.* **137**, 11461–11475 (2015).
- Row, R. D. & Prescher, J. A. Constructing new bioorthogonal reagents and reactions. *Acc. Chem. Res.* **51**, 1073–1081 (2018).
- Kumar, A., Li, K. & Cai, C. Anaerobic conditions to reduce oxidation of proteins and to accelerate the copper-catalyzed “click” reaction with a water-soluble bis(triazole) ligand. *Chem. Commun.* **47**, 3186–3188 (2011).
- Egorova, K. S. & Ananikov, V. P. Which metals are green for catalysis? comparison of the toxicities of Ni, Cu, Fe, Pd, Pt, Rh, and Au salts. *Angew. Chem. Int. Ed.* **55**, 12150–12162 (2016).
- Plutschack, M. B., Pieber, B., Gilmore, K. & Seeberger, P. H. The Hitchhiker’s guide to flow chemistry. *Chem. Rev.* **117**, 11796–11893 (2017).
- Noël T., Su Y., Hessel V. in *Organometallic Flow Chemistry*, (eds Noël T.) Vol. 57. (Switzerland, Springer, 2015).
- Barnes, J. C. et al. Iterative exponential growth of stereo- and sequence-controlled polymers. *Nat. Chem.* **7**, 810–815 (2015).
- Leibfarth, F. A., Johnson, J. A. & Jamison, T. F. Scalable synthesis of sequence-defined, unimolecular macromolecules by flow-IEG. *Proc. Natl Acad. Sci. USA* **112**, 10617–10622 (2015).
- Mijalis, A. J. et al. A fully automated flow-based approach for accelerated peptide synthesis. *Nat. Chem. Biol.* **13**, 464–466 (2017).
- Bogdan, A. R. & Sach, N. W. The use of copper flow reactor technology for the continuous synthesis of 1,4-disubstituted 1,2,3-triazoles. *Adv. Synth. Catal.* **351**, 849–854 (2009).
- Fuchs, M., Goessler, W., Pilger, C. & Kappe, C. O. Mechanistic insights into copper(I)-catalyzed azide–alkyne cycloadditions using continuous flow conditions. *Adv. Synth. Catal.* **352**, 323–328 (2010).
- Razzaq, T. & Kappe, C. O. Continuous flow organic synthesis under high-temperature/pressure conditions. *Chem. Asian J.* **5**, 1274–1289 (2010).
- Bogdan, A. R. & James, K. Efficient access to new chemical space through flow—construction of druglike macrocycles through copper-surface-catalyzed azide–alkyne cycloaddition reactions. *Chem. Eur. J.* **16**, 14506–14512 (2010).
- Smith, C. J., Nikbin, N., Ley, S. V., Lange, H. & Baxendale, I. R. A fully automated, multistep flow synthesis of 5-amino-4-cyano-1,2,3-triazoles. *Org. Biomol. Chem.* **9**, 1938–1947 (2011).
- Carlos Varas, A., Noël, T., Wang, Q. & Hessel, V. Copper(I)-catalyzed azide–alkyne cycloadditions in microflow: catalyst activity, high-t operation, and an integrated continuous copper scavenging unit. *ChemSusChem* **5**, 1703–1707 (2012).
- Ötvös, S., Mándity, I. M., Kiss, L. & Fülöp, F. Alkyne–azide cycloadditions with copper powder in a high-pressure continuous-flow reactor: high-temperature conditions versus the role of additives. *Chem. Asian J.* **8**, 800–808 (2013).
- Ötvös, S., Georgiades, A., Mándity, I. M., Kiss, L. & Fülöp, F. Efficient continuous-flow synthesis of novel 1,2,3-triazole-substituted β -aminocyclohexanecarboxylic acid derivatives with gram-scale production. *Beilstein J. Org. Chem.* **9**, 1508–1516 (2013).
- Ötvös, S. et al. Continuous-flow azide–alkyne cycloadditions with an effective bimetallic catalyst and a simple scavenger system. *RSC Adv.* **4**, 46666–46674 (2014).
- Gürsel, I. V., Aldiansyah, F., Wang, Q., Noël, T. & Hessel, V. Continuous metal scavenging and coupling to one-pot copper-catalyzed azide–alkyne cycloaddition click reaction in flow. *Chem. Eng. J.* **270**, 468–475 (2015).
- Chassaing, S., Beénéteau, V. & Pale, P. When CuAAC ‘Click Chemistry’ goes heterogeneous. *Catal. Sci. Technol.* **6**, 923–957 (2016).
- Mandoli, A. Recent advances in recoverable systems for the copper-catalyzed azide–alkyne cycloaddition reaction (CuAAC). *Molecules* **21**, 1174 (2016).
- Seath, C. P., Burley, G. A. & Watson, A. J. B. Determining the origin of rate-independent chemoselectivity in CuAAC reaction: an alkyne-specific shift in rate-determining step. *Angew. Chem. Int. Ed.* **56**, 3314–3318 (2017).
- Hatit, M. Z. C. et al. Chemoselective Sequential Click Ligations Directed by Enhanced Reactivity of an Aromatic Ynamine. *Org. Lett.* **18**, 1694–1697 (2016).
- Reiner, T., Earley, S., Turetsky, A. & Weissleder, R. Bioorthogonal small-molecule ligands for PARP1 imaging in living cells. *ChemBiochem* **11**, 2374–2377 (2010).
- O’Donovan, L. et al. Parallel synthesis of cell-penetrating peptide conjugates of PMO toward exon skipping enhancement in duchenne muscular dystrophy. *Nucleic Acid. Ther.* **25**, 1–10 (2015).
- Re, F. et al. Functionalization of liposomes with ApoE-derived peptides at different density affects cellular uptake and drug transport across a blood-brain barrier model. *Nanomedicine* **7**, 551–559 (2011).
- Godfrey, C. et al. Delivery is key: lessons learnt from developing splice-switching antisense therapies. *EMBO Mol. Med.* **9**, 545–557 (2017).
- Sharma, V. K., Sharmab, R. K. & Singh, S. K. Antisense oligonucleotides: modifications and clinical trials. *Med. Chem. Commun.* **5**, 1454–1471 (2014).
- Juliano, R. L. The delivery of therapeutic oligonucleotides. *Nucleic Acids Res.* **44**, 6518–6548 (2016).
- Shabanpoor, F. et al. Identification of a peptide for systemic brain delivery of a morpholino oligonucleotide in mouse models of spinal muscular atrophy. *Nucleic Acid Ther.* **27**, 130–143 (2017).

Acknowledgements

We thank Vapourtec for the copper reactors, the EPSRC UK National Mass Spectrometry Facility at Swansea University for analyses, the University of Strathclyde for PhD studentship (M.Z.C.H.), the Leverhulme Trust for a postdoctoral fellowship (L.F.R.; grant no. RPG-2014–001), and the BBSRC for funding (grant nos. BB/R006857/1, BB/N016378/1).

Author contributions

Conceptualization: G.A.B., A.J.B.W.; data analysis: M.Z.C.H., G.A.B., A.J.B.W.; synthetic chemistry: M.Z.C.H., J.M.T.; preparation of peptide-based materials: M.Z.C.H.;

preparation of phosphoramidite building blocks and solid phase ODN synthesis: L.F.R.; project supervision: F.V., G.A.B, A.J.B.W.; writing of the paper: G.A.B, A.J.B.W.

Additional information

Supplementary Information accompanies this paper at <https://doi.org/10.1038/s41467-018-06551-0>.

Competing interests: The authors declare no competing interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018